

AD-A107 950

CALIFORNIA UNIV LOS ANGELES DEPT OF CHEMISTRY

F/G 20/2

SPECTRAL DIFFUSION IN ORIENTATIONALLY DISORDERED ORGANIC SOLIDS--ETC(U)

NOV 81 P N PRASAD, J R MORGAN, M A EL-SAYED

N00014-75-C-0602

NL


UNCLASSIFIED

TR-39

ISI
SERIAL



END
DATE
FILMED
BY
PTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

LEVEL II

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0602

Task No. NR 056-498

TECHNICAL REPORT NO. 39

SPECTRAL DIFFUSION IN ORIENTATIONALLY
DISORDERED ORGANIC SOLIDS

by

Paras N. Prasad, Jack R. Morgan and Mostafa A. El-Sayed
Department of Chemistry
University of California
Los Angeles, California 90024

DTIC

DEC 01 1981

Accepted for Publication

in

The Journal of Physical Chemistry

November 20, 1981

Reproduction in whole or in part is permitted for any
purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

81 11 30 067

AD A107950

DTIC FILE COPY

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. 39	2. GOVT ACCESSION NO. AD-A107950	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Spectral Diffusion in Orientationally Disordered Organic Solids		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Paras N. Prasad, Jack R. Morgan and Mostafa A. El-Sayed		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0602
9. PERFORMING ORGANIZATION NAME AND ADDRESS Regents of the University of California University of California, 405 Hilgard Ave. Los Angeles, CA 90024		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-056-498
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Branch Arlington, Virginia 22217		12. REPORT DATE November 20, 1981
		13. NUMBER OF PAGES 16
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Office of Naval Research Branch Office 1030 East Green Street Pasadena, CA 91106		15. SECURITY CLASS. (of this report) unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; distribution of this document is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Accepted for publication in J. Phys. Chem.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Spectral diffusion Luminescence Line narrowing Orientational disorder		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The spectral diffusion within the inhomogeneous linewidth of the $T_1 \rightarrow S_0$ transition is presented for an orientationally disordered organic solid under the condition when kT ($\sim 3 \text{ cm}^{-1}$) is much smaller than the inhomogeneous linewidth ($\sim 83 \text{ cm}^{-1}$). The rate of the transfer is found to increase as the site energy increases. This result is explained in terms of a phonon assisted unidirectional (high to low) energy transfer process in which the increasing density of the low energy sites (acceptors) with an increase in the excitation energy plays an important role.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SPECTRAL DIFFUSION IN ORIENTATIONALLY
DISORDERED ORGANIC SOLIDS

Paras N. Prasad⁺, Jack R. Morgan and Mostafa A. El-Sayed
Chemistry Department
University of California
Los Angeles, CA 90024

ABSTRACT

The spectral diffusion within the inhomogeneous linewidth of the $T_1 \leftarrow S_0$ transition is presented for an orientationally disordered organic solid under the condition when kT ($\sim 3 \text{ cm}^{-1}$) is much smaller than the inhomogeneous linewidth ($\sim 83 \text{ cm}^{-1}$). The rate of the transfer is found to increase as the site energy increases. This result is explained in terms of a phonon assisted unidirectional (high to low) energy transfer process in which the increasing density of the low energy sites (acceptors) with an increase in the excitation energy plays an important role.

⁺Alfred P. Sloan Fellow Permanent Address: The Department of Chemistry, State University of New York at Buffalo, Buffalo, New York 14914

Accession For	
NTIS Grant	<input checked="" type="checkbox"/>
DTIC	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution _____	
Availability _____	
Dist	<input type="checkbox"/>
A	

INTRODUCTION

Great interest has recently developed in the optical properties of, and energy transfer in, amorphous solids, e.g., inorganic glasses. Both the inhomogeneous and homogeneous linewidths for T less than $\sim 100\text{K}$ of electronic transitions of impurity ions in these solids are found to be broader than in crystalline solids^{1,2}. Fluorescence line narrowing techniques³ have recently been used to study the spectral diffusion of the $^5\text{D}_0 \leftarrow ^7\text{F}_0$ transition energy of Eu^{3+} amongst different sites in calcium metaphosphate glass at temperatures ($200^\circ - 350^\circ \text{K}$) where kT is greater than the inhomogeneous width ($\sim 50 \text{ cm}^{-1}$). The results suggest⁴ that the transfer is dipolar in nature, phonon assisted, and is independent of the donor site-energy within the inhomogeneous profile. Indications are that the diffusion limit (i.e., the limit when the spectral diffusion coefficient is independent of time) is reached only when the excitation level of the donor is reduced to 5% of its initial value at room temperature and 30 mole percent Eu^{3+} concentration⁵.

Orientationally disordered molecular solids are a group of solids which retain a high degree of translation correlation in terms of lattice positions of the center of mass but are disordered generally with respect to the position of the substituent groups. Thus, they represent a class inbetween that of an ordered crystalline solid and a truly amorphous medium (it might thus be called an amorphalline solid!) These solids give rise to well defined diffraction patterns that yield a higher average crystal symmetry. The orientational disorder leads to a profound broadening of the spectral profile of the electronic

transitions⁶. The broadening generally results from an inhomogeneous distribution of the site energy (i.e., a variation in the D-shift term in the language of the Frenkel exciton theory)⁷.

For understanding the nature of spectral diffusion in a disordered system, these groups of solids are a more useful protosystem than other disordered solids having the following distinct merits:

(i) As the disorder is inherent within the average higher crystal symmetry, it is found to be independent of the history of the crystal preparation. Thus, the observed spectral profile appears to be insensitive to the sample history and is highly reproducible. On the other hand, inhomogeneities in crystals and disorder structure in a glass are very much dependent on the method of the sample preparation creating a problem of reproducibility of spectral profile.

(ii) Often, there is an analog ordered system with which a comparison can be made to discern the effect of disorder both on the electronic states and the phonon bands. The results of investigation of phonon bands indicates that the presence of orientational disorder does not effectively destroy the k-symmetry of the phonon states and thus, no phonon localization is observed. On the other hand, for the case of spectral diffusion for an impurity in a glassy system⁴, the possibility of phonon localization at the impurity site becomes a complicating factor in the interpretation of the result.

The present study focuses on a specific system, 1-bromo-4 chloronaphthalene (from here on abbreviated as BCN) which exhibits disorder with respect to the position of the halogens⁸. The observed linewidth⁶ of the $T_1 \leftarrow S_0$ absorption of this crystal at 4.2°K is 83 cm^{-1}

(i.e., kT is \ll the linewidth). This linewidth is about two orders of magnitude larger than the linewidth observed ($< 1 \text{ cm}^{-1}$) for the corresponding transition in 1, 4 - dichloronaphthalene⁶ (DCN) which crystalizes in the same space group with an ordered structure⁸, suggesting that the width in BCN is broadened by the orientational disorder. Although the optical phonon transitions observed in the Raman spectra of BCN are found to be broader than that observed for DCN, an excellent phonon spectral correlation⁸ between the two crystals reveals that the k-symmetry selection rule for the disordered BCN is not completely destroyed for the optical phonons.

In addition to the fact that BCN is a different type of disordered system than glasses and that the $T_1 \leftarrow S_0$ transition has a linewidth which is much larger than kT , the mechanism of the transfer of the $T_1 \leftarrow S_0$ transition energy is of the exchange type (not dipolar in nature as in the system studied previously^{4,5}). The detailed comparison of the nature of the energy diffusion in the two systems would be most important.

EXPERIMENTAL

The BCN was synthesized from 1-amino-4-chloronaphthalene using the procedure described elsewhere⁴. The material was extensively zone refined. Crystals were grown from the melt in a Bridgmann furnace. Samples of 0.5 x 0.5 x 0.2 cm size were mounted in a liquid helium dewar. A Quanta-Ray DCR-1 Nd:YAG pumped pulsed dye laser with a spectral width of $\sim 0.3 \text{ cm}^{-1}$ and a pulse width of 6 n sec was used as the $S_0 \rightarrow T_1$ excitation source. The emission was dispersed using a

1 meter Jarrell-Ash monochromator with a 2 cm^{-1} slit width. Time resolved spectra of the first vibronic band of the $T_1 - S_0$ phosphorescence were recorded using a PAR model 162 boxcar averager. A special gated phototube was used in order to reject scattered laser light.

RESULTS

The relationship between the absorption spectra and the emission spectra of the triplet state in 1-bromo-4-chloronaphthalene has been reported elsewhere⁶. It is to be noted that the emission in a steady state experiment is observed from the low energy sites. Furthermore, the width of the emission is about 20 cm^{-1} which is much narrower than the absorption linewidth but is still considerably broader than that observed for the ordered 1,4 dichloronaphthalene triplet emission.

Figure 1 demonstrates that spectral diffusion of the $T_1 \rightarrow S_0$ transition energy occurs in this orientationally disordered material. The band shape of the first vibronic band was monitored as a function of delay time. The spectra displayed were obtained by exciting a site on the low energy side of the $S_0 \rightarrow T_1$ absorption profile. The figure shows that as the delay time increases the resonant type emission decreases in intensity while emission from lower energy acceptors increases in intensity. This figure also shows the same spectra recorded at two different sample temperatures. The phosphorescence spectra obtained under identical conditions but at two different sample temperatures, clearly indicate that the transfer rate is temperature dependent, and thus, phonon assisted⁹.

The results of the effect of site selective excitation on the rate of spectral diffusion of the $T_1 \rightarrow S_0$ energy is shown in fig. 2. In each case the spectrum is sampled after a 10 μsec delay using a 50 μsec sampling time. In order to minimize interference with stray light due to the laser,

the first vibronic band profile of the emission is monitored. It can be seen that the observed emission profile is very much dependent on the excitation energy. As the low energy sites are excited, a relatively narrow resonant emission is observed, indicating a transfer rate slower than the sampling time. On the other hand with excitation of higher energy sites, the emission profile is broadened indicating a faster transfer time to a broader distribution of the low energy sites. These observations are suggestive of a transfer rate which at 4.2 K is dependent on the donor site energy.

DISCUSSION

The observed emission line narrowing with the excitation of the low energy sites clearly indicates that the singlet-triplet absorption in BCN is inhomogeneously broadened. The efficient energy transfer from the highest energy sites to the lowest energy sites, without the observation of mobility edges, might suggest the absence of Anderson type localization in this disordered system.

In our previous studies⁴ on the transfer of Eu^{3+} excitation in phosphate glasses at room temperature, we found that the full inhomogeneous profile was recovered uniformly with time following pulsed laser excitation, independent of the laser wavelength used. This suggests that the transfer rate is independent of donor-acceptor energy mismatch. This is expected to be the case when $kT \gg$ the inhomogeneous width. In our present system, however, kT is $\sim 3 \text{ cm}^{-1}$ and the inhomogeneous width is $\sim 80 \text{ cm}^{-1}$. Transfer is thus only possible from high to low energy sites at this temperature. This is because transfer from a low energy site to a high energy site requires the population of phonons which is very small at 4.2°K. For the same reason,

the Raman and Orbach type phonon assisted processes⁹ would also be unimportant at this low temperature, thus leaving the direct one phonon emission process as the most probable mechanism for the observed spectral diffusion in this system. For this process, the transfer rate from a donor at site energy E_i can be written as:

$$W(E_i) \propto \sum_f c_{if, \omega_{if}} \rho_f(E_i) \rho(\omega_{if}) (n_{\omega_{if}} + 1) \quad (1)$$

In this equation c_{if} contains the details of the coupling with phonons required to make up the energy mismatch between the energy of the donor and that of the acceptor; $\rho_f(E_i)$ is the density of the acceptor states at energy $\leq E_i$ to which energy can be transferred at 4.2⁰K by the emission of one phonon. (the inhomogeneous profile can be assumed to result from a distribution of sites). The term n_{if} represents the occupation number of phonons of frequency $\omega_{if} = \frac{E_i - E_f}{h}$ and $\rho(\omega_{if})$ is the phonon density of states. At 4.2⁰K, $(n_{\omega_{if}} + 1)$ is near unity.

Both for small and large energy mismatch, $c_{if, \omega_{if}} \times \rho(\omega_{if})$ increases with the increase of the energy mismatch (i.e., increase in donor site energy). Furthermore, as the energy of the donor, E_i , increases, the density of the acceptors increases (which leads to an increase in the phase space for the energy transfer process). Thus as a result of all these terms, an increase in the transfer rate is expected as the donor energy increases. This fact explains the lack of emission from sites except the ones at the lowest energy. It also explains our spectral diffusion results which show an increase of the emission from the low energy sites with time upon exiting the higher energy donors.

The observations of the present work are summarized as follows:

(i) The spectral profile for the singlet-triplet transition is inhomogeneously broadened by the orientational disorder, (ii) No Anderson type localization is observed for the triplet excitation in this disordered system (iii) The energy transfer is phonon assisted (iv) The spectral diffusion rate is site dependent and increases with an increase in the donor energy.

A more quantitative correlation of the theory and the results will require quantitative measurements of the transfer rates of various low energy sites. Such measurements are in progress and will be reported at a later date.


Acknowledgements: P.P. wishes to thank Petroleum Research Fund, J.M. and M.A.E. thank the Office of Naval Research for financial support of this work.


REFERENCES

- 1.) P.M. Selzer, D.L. Huber, D.S. Hamilton, W.M. Yen and M.J. Weber, Phys. Rev. Lett. 36, 813(1976).
- 2.) J. Hegarty and W.M. Yen, Phys. Rev. Lett. 43, 1126(1976).
- 3.) A. Szabo, Phys. Rev. Lett. 25, 924(1970).
- 4.) P. Avouris, A. Campion and M.A. El-Sayed in "Advances in Laser Spectroscopy I", p. 57.
- 5.) J.R. Morgan and M.A. El-Sayed, J. Phys. Chem., to be published.
- 6.) J.C. Bellows and P.N. Prasad, to be published;
L.A. Hess and P.N. Prasad, unpublished results.
- 7.) D.P. Craig and S.H. Walmsley, Exciton in Molecular Crystals, W.A. Benjamin, Amsterdam, 1968.
- 8.) J.C. Bellows and P.N. Prasad, J. Chem. Phys. 67, 5802(1979);
J.C. Bellows, E.D. Stevens, and P.N. Prasad, Acta Cryst. 1334, 3256(1978).
- 9.) T. Holstein, S.K. Lyo and R. Orbach, Phys. Rev. Lett. 36, 891(1976).

Fig. 1. Spectral diffusion of singlet-triplet excitation energy in an "amorphalline" solid at 4.2°K using time resolved phosphorescence line narrowing techniques. The system is 1-bromo-4-chloronaphthalene excited at 4943 Å. The top three spectra are of the $0,0 - 321\text{ cm}^{-1}$ vibronic band of the phosphorescence emission recorded at different delay times after the laser pulsed excitation with a 50 μsec sampling time. The bottom spectrum illustrates the effect of temperature on the rate of spectral diffusion within the inhomogeneous profile of the $0,0 - 321\text{ cm}^{-1}$ band.

Fig. 2. Dependence of the spectral diffusion rate of the singlet-triplet excitation on the donor site energy in 1-bromo-4-chloronaphthalene at 4.2°K. The spectral intensity change of the $0,0 - 321\text{ cm}^{-1}$ band of the $T_1 \rightarrow S_0$ phosphorescence is monitored with a 10 μsec delay using a 50 μsec sampling time following pulsed laser excitation at different wavelength (i.e., different donor energies) within the inhomogeneously broadened $0,0$ band of the $T_1 \leftarrow S_0$ absorption. The results show that the ratio of the line narrowed (donor) phosphorescence to the lower energy broad emitting traps decreases as the donor (excitation) energy increases, suggesting an increase in the spectral diffusion rate.





TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Western Regional Office Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Eastern Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
		Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>
Mr. James Kelley DTNSRDC Code 2803 Annapolis, Maryland 21402	1
Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. M. A. El-Sayed Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1	Dr. M. Rauhut Chemical Research Division American Cyanamid Company Bound Brook, New Jersey 08805	1
Dr. E. R. Bernstein Department of Chemistry Colorado State University Fort Collins, Colorado 80521	1	Dr. J. I. Zink Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. D. Haarer IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. J. R. MacDonald Chemistry Division Naval Research Laboratory Code 6110 Washington, D.C. 20375	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. G. B. Schuster Chemistry Department University of Illinois Urbana, Illinois 61801	1	Dr. William M. Jackson Department of Chemistry Howard University Washington, DC 20059	1
Dr. A. Adamson Department of Chemistry University of Southern California Los Angeles, California 90007	1	Dr. George E. Walraffen Department of Chemistry Howard University Washington, DC 20059	1
Dr. M. S. Wrighton Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1		

